## New Jersey Division of AIDS Prevention and Control Persons Living with HIV/AIDS as of June 30, 2002 Mercer County Residents

			Adult/A	Adol	escei	nt *		Pediat	ric *			Tota	al	
1.	Current HIV/AIDS (		Cases		% 			Cases	% 			lases	% 	
	AIDS-Any OI AIDS-CD4 Only HIV Only		137 352 508	7 2 8	14 35 51			8 0 7	53 ( 47	3 ) 7		145 352 515	1 3 5	4 5 1
	Total		995		100			15	100			1012	10	
2.	Current Age * Cases %		3. Rac					Adult/ Cases	8 %	Cases	8	Cases	%	
	<pre>&lt;5  # # 5-12  # # 13-19  9  1 20-29  58  6</pre>		Whi Bla His Oth	ite, ack, span her/	Not Not ic Unkno	own	anic anic	 # 695 # 9	# 70 # 1	# 0	# 80 # 0	104 9	19 70 10 1	
	30-39 322 32 40-49 395 39 50+ 213 21 Unk. 0 0 Total 1012 100			tal				997		15		1012		
4.	Patient Group				Ma:	Ad les		dolescer	nt Tran Female		-	Iodes ** Tota	al	
	Men sex with men ( Injection drug use MSM/IDU Heterosexual conta Risk not reported/	MSM) er (IDU) act other***				70 138	30 32 5 11 22		126 ( 13! 111	0 0 5 34 0 0 5 36 1 30		32 3 20 24	35 21 30 05 49	1 3 2 2
	Total					618	100		372	2 100		99	90	10
						les	%	diatric	Female	es %		Tota		%
	Parent at risk/has	S AIDS/HIV other***				8 #	80 20			9 75 ‡ 25	; ;	1	L7 5	 7 2

NOTE: Includes only those persons known to be infected with HIV.

Total

- # Indicates that number is not shown due to small cell size, in accordance with NJDHSS security and confidentiality policies.
- \* Classification based on current age at the end of June 2002.
- \*\* 7 patients are currently of adult/adolescent age (13+ years old), but have evidence of having been HIV-infected as children. They are counted as adult/adolescent reports in tables 1,2, and 3, and as pediatric reports in table 4.

10 100

12 100

22 100

\*\*\* Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

#### New Jersey Division of AIDS Prevention and Control AIDS Cases Reported as of June 30, 2002 Mercer County Residents

Adult/Adolescent\*

Cases (%)

1. Disease Category

Total

PCP Other OI w KS Alone No Disease			# ( #) # ( #) 15 ( 1) 576 ( 48)	# ( #) # ( #) 12 ( 80) 224 ( 39)	# 0	( #) ( #) ( 0)	# C	# ( #) # ( #) ) ( .)	•	28)	244 257 12 224	( 7
Total		1	211 (100)	724 ( 60)	23	(100)	13	3 ( 57)	1234 (1	.00)	737	( 6
Age *	Cases	%	3. Race/E	th.	Adult/A Cases		Pediatr Cases	cic*	Total Cases	%		
Under 5	#	#	White,	Not Hispanic	#	#	#	#	297	24		
5-12	#	#	Black,	Not Hispanic	796	66	20	87	816	66		
13-19	6	0	Hispan	ic	#	#	#	#	113	9		
20-29	150	12	Other/	Unknown	8	1	0	0	8	1		
30-39	503	41										
40-49	394	32	Total		1211	100	23	100	1234	100		
Over 49	158	13										
Unknown	0	0										
Total	1234	100										

Deaths (%) Cases (%)

Pediatric\*

Deaths (%) Cases (%)

Total

Deaths (%)

Risk not reported/other***	0	0	0	0	0	0			
Parent at risk/has AIDS/HIV	12	100	11	100	23	100			
	Males	%	Females	8	Total	왕			
	Pediatric Transmission Modes								
Total	877	100	334	100	1211	100			
Risk not reported/other***	122	14	79	24	201	17			
Heterosexual contact	71	8	97	29	168	14			
MSM/IDU	54	6	0	0	54	4			
Injection drug user (IDU)	354	40	158	47	512	42			
Men sex with men (MSM)	276	31	0	0	276	23			
	Males	%	Females	%	Total	왕			
Patient Group	Adult/Adolescent Transmission Modes **								

12 100

11 100

23 100

<sup>#</sup> Indicates that number is not shown due to small cell size, in accordance with NJDHSS security and confidentiality policies.

<sup>\*</sup> Classification at time of AIDS diagnosis.

<sup>\*\* 0</sup> patients were diagnosed with AIDS as adults, but have evidence either of having been HIV-infected or of having acquired AIDS as children. They are counted as adult/adolescent reports in tables 1,2, and 3; and as pediatric reports in table 4.

<sup>\*\*\*</sup> Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

### New Jersey Division of AIDS Prevention and Control AIDS Cases Reported as of June 30, 2002 Mercer County Residents

5. AIDS Cases by Year of Diagnosis.

Year of Diagnosis	Number of Cases	Deaths	Case-Fatality Rate
Before 1985	13	13	100%
1985	15	15	100%
1986	15	15	100%
1987	33	33	100%
1988	38	35	92%
1989	52	46	88%
1990	45	44	98%
1991	73	68	93%
1992	108	91	84%
1993	153	106	69%
1994	117	76	65%
1995	133	76	57%
1996	95	34	36%
1997	74	21	28%
1998	58	19	33%
1999	62	14	23%
2000	74	18	24%
2001	63	#	#%
2002	13	#	#%
Total	1234	737	60%

<sup>#</sup> Indicates that number is not shown due to small cell size, in accordance with NJDHSS security and confidentiality policies.

### New Jersey Division of AIDS Prevention and Control HIV INFECTION-not-AIDS Reports with Identifiers Reported as of June 30, 2002 Mercer County Residents

1.	Age *	Cases	%	2.	Race/Ethnicity	Adult/ Cases		Pedia Cases			Total ases	%	
	Under 5	#	#		White, Not Hispanic	103	18	6	50		109	19	
	5-12	#	#		Black, Not Hispanic	413	72	5	42		418	71	
	13-19	13	2		Hispanic	#	#	#	#		52	9	
	20-29	149	25		Other/Unknown	#	#	#	#		9	2	
	30-39	212	36							_			
	40-49	136	23		Total	576	100	12	100		588	100	
	Over 49	66	11										
	Unknown	0	0										
	Total	588	100										
3.	Patient Gro	up			Adult/Adolescent Transmission Modes **								
					Males	%	Fer	males %		Total	%		
	Men sex wit	h men (M	SM)		99	29		0 0		99	17		
	Injection d				104	30		83 36		187	32		
	MSM/IDU		, _ ,		15	4		0 0		15	3		
	Heterosexua	l contac	t		42	12		82 36		124	22		
	Risk not re			*	86	25		65 28		151	26		

	Pediatric Transmission Modes						
	Males	%	Females %	Total	%		
Parent at risk/has AIDS/HIV	#	#	# #	7	58		
Risk not reported/other***	#	#	# #	5	42		
Total	6	100	6 100	12	100		

\_\_\_\_

346 100

-----

230 100

576 100

\_\_\_\_\_

NOTE: Includes only those persons known to be infected with HIV.

Total

- Indicates that number is not shown due to small cell size, in accordance with NJDHSS security and confidentiality policies.
- Classification at time of first HIV+ test or at time of report if not yet confirmed HIV+.
- \*\* O patients were diagnosed with HIV as adults, but have evidence of having been HIV-infected as children. They are counted as adult/adolescent reports in tables 1 and 2; and as pediatric reports in table 3.
- \*\*\* Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

NOTE: These tables include those reports of HIV Positive infection that have not yet been updated to AIDS status. Not included in these tables are cases initially reported as HIV-not-AIDS that have subsequently been updated to AIDS. Also not included in these tables are HIV Perinatal Exposure Reports where the serostatus of the child is not confirmed HIV Positive; i.e., the HIV serostatus of the child is undetermined or has been found to be negative (seroreverted).

### New Jersey Division of AIDS Prevention and Control HIV/AIDS Cases Reported as of June 30, 2002 Mercer County Residents

						Pediatric *				Total					
1.	Current HI				Cases		% 	Case		왕 		Cas		) 	
	AIDS-Any O AIDS-CD4 O HIV Only	oI Only			635 576 576		36 32	:	23 0	66 0 34		6 5	558 576 588	36 32	
	Total				1787		100		35	100				100	
2.	Age *			3.				Adult/A	%	Cas		%		To Cases	tal %
	Under 5 5-12 13-19 20-29 30-39 40-49 Over 49 Unknown	#	# 1 19 40 27		White, Black, Hispani Other/U	Not Not c nkn	Hispanic Hispanic own	398	22 68 # 		8 25 #	23 71 # #		406 1234 165 17  1822	9
	Total	1822	100												
4.	Patient Gr	-					Males	ılt/Adole: %		Transm		-			
	Men sex wi Injection MSM/IDU Heterosexu Risk not r	drug user al contac eported/c	MSM) (IDU) ct other***				375 458 69 113 208	37 6 9	_	0 241 0 179 144	43 0 32 26		375 699 69 292 352	39 4 16 20	

1787 100

564 100

	Pediatric Transmission Modes						
	Males	%	Females	%	Total	%	
Parent at risk/has AIDS/HIV	16	89	14	82	30	86	
Risk not reported/other***	#	11	#	18	5	14	
Total	18	100	17	100	35	100	

1223 100

NOTE: Includes only those persons known to be infected with HIV.

- Indicates that number is not shown due to small cell size, in accordance with NJDHSS security and confidentiality policies.
- \* Classification at time of first HIV+/AIDS diagnosis.

Total

- \*\* 0 patients were diagnosed with HIV/AIDS as adults, but have evidence of having been HIV-infected as children. They are counted as adult/adolescent reports in tables 1,2 and 3; and as pediatric reports in table 4.
- \*\*\* Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

# New Jersey Division of AIDS Prevention and Control HIV/AIDS Cases Reported as of June 30, 2002 Mercer County Residents

5. HIV/AIDS Cases by Year of First HIV/AIDS Diagnosis.

Year of First HIV/AIDS Diagnosis	Cases	Number Known Dead	Case-Fatality Rate
Before 1985	17	14	82%
1985	17	17	100%
1986	27	24	89%
1987	46	43	93%
1988	48	40	83%
1989	80	66	83%
1990	93	74	80%
1991	111	73	66%
1992	232	129	56%
1993	185	82	44%
1994	188	83	44%
1995	175	56	32%
1996	139	36	26%
1997	101	18	18%
1998	95	22	23%
1999	96	15	16%
2000	96	12	13%
2001	58	6	10%
2002	17	0	0%
Unknown	#	0	0%
Total	1822	810	44%

<sup>#</sup> Indicates that number is not shown due to small cell size, in accordance with NJDHSS security and confidentiality policies.